



Staphylococcus aureus Coproporphyrinogen III Oxidase Is Required for Aerobic and Anaerobic Heme Synthesis

 Jacob E. Choby,^{a,b,*} Eric P. Skaar^{a,c}

^aDepartment of Pathology, Microbiology, and Immunology, Vanderbilt University Medical Center, Nashville, Tennessee, USA

^bGraduate Program in Microbiology and Immunology, Vanderbilt University, Nashville, Tennessee, USA

^cVanderbilt Institute for Infection, Immunology, and Inflammation, Vanderbilt University Medical Center, Nashville, Tennessee, USA

ABSTRACT The virulence of the human pathogen *Staphylococcus aureus* is supported by many heme-dependent proteins, including key enzymes of cellular respiration. Therefore, synthesis of heme is a critical component of staphylococcal physiology. *S. aureus* generates heme via the coproporphyrin-dependent pathway, conserved across members of the *Firmicutes* and *Actinobacteria*. In this work, we genetically investigate the oxidation of coproporphyrinogen to coproporphyrin in this heme synthesis pathway. The coproporphyrinogen III oxidase CgoX has previously been identified as the oxygen-dependent enzyme responsible for this conversion under aerobic conditions. However, because *S. aureus* uses heme during anaerobic nitrate respiration, we hypothesized that coproporphyrin production is able to proceed in the absence of oxygen. Therefore, we tested the contribution to anaerobic heme synthesis of CgoX and two other proteins previously identified as potential oxygen-independent coproporphyrinogen dehydrogenases, NWMN_1486 and NWMN_1636. We have found that CgoX alone is responsible for aerobic and anaerobic coproporphyrin synthesis from coproporphyrinogen and is required for aerobic and anaerobic heme-dependent growth. This work provides an explanation for how *S. aureus* heme synthesis proceeds under both aerobic and anaerobic conditions.

IMPORTANCE Heme is a critical molecule required for aerobic and anaerobic respiration by organisms across kingdoms. The human pathogen *Staphylococcus aureus* has served as a model organism for the study of heme synthesis and heme-dependent physiology and, like many species of the phyla *Firmicutes* and *Actinobacteria*, generates heme through a coproporphyrin intermediate. A critical step in terminal heme synthesis is the production of coproporphyrin by the CgoX enzyme, which was presumed to be oxygen dependent. However, *S. aureus* also requires heme during anaerobic growth; therefore, the synthesis of coproporphyrin by an oxygen-independent mechanism is required. Here, we identify CgoX as the enzyme performing the oxygen-dependent and -independent synthesis of coproporphyrin from coproporphyrinogen, resolving a key outstanding question in the coproporphyrin-dependent heme synthesis pathway.

KEYWORDS *Staphylococcus*, heme, tetrapyrroles

Staphylococcus aureus is an important human pathogen responsible for a wide variety of diseases in numerous host niches (1) and is supported by a robust host-adapted metabolism. *S. aureus* is a facultative anaerobe of the *Firmicutes* phylum capable of performing oxygen-dependent cellular respiration or transitioning to anaerobic respiration in the presence of alternative terminal electron acceptors like nitrate. In the absence of respiration, fermentation of a variety of carbon sources supports *S. aureus* replication. The tetrapyrrole cofactor heme is important for the cellular processes of many organisms across kingdoms, and in *S. aureus*, heme is required for aerobic and anaerobic respiration. Heme is a cofactor of the succinate dehydrogenase

Citation Choby JE, Skaar EP. 2019.


Staphylococcus aureus coproporphyrinogen III oxidase is required for aerobic and anaerobic heme synthesis. mSphere 4:e00235-19. <https://doi.org/10.1128/mSphere.00235-19>.

Editor Paul D. Fey, University of Nebraska Medical Center

Copyright © 2019 Choby and Skaar. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Eric P. Skaar, eric.skaar@vumc.org.

* Present address: Jacob E. Choby, Emory Vaccine Center and Emory Antibiotic Resistance Center, Emory University School of Medicine, Atlanta, Georgia, USA.

 Anaerobic heme synthesis in *S. aureus* @jake_choby @EricSkaar

Received 3 April 2019

Accepted 26 June 2019

Published 10 July 2019

(SdhC; cytochrome b_{558} subunit [2, 3]) that reduces menaquinone, the two terminal cytochrome oxidases (QoxABCD and CydAB [4, 5]) that oxidize menaquinol during aerobic respiration, and the nitrate reductase (NarI; gamma subunit [6, 7]) that oxidizes menaquinol during anaerobic nitrate respiration. Therefore, mutants unable to synthesize heme are respiration deficient and grow poorly, adopting a “small-colony variant” phenotype most evident on solid medium (8). Additionally, *S. aureus* and other species rely on heme for the function of many widely conserved enzymes, including catalase, nitric oxide synthase, and globin family proteins.

Bacteria synthesize heme from the universal precursor δ -aminolevulinic acid (ALA) which proceeds to uroporphyrinogen. Uroporphyrinogen can be diverted to form similar cofactors, including siroheme for use in the *S. aureus* nitrite reductase (9). Bacteria use different pathways to synthesize heme from uroporphyrinogen, which has been extensively reviewed by Dailey and colleagues (10). In only the last decade, it has been recognized that species of *Firmicutes* and *Actinobacteria* synthesize heme via a unique pathway that uses the coproporphyrin intermediate (11, 12). The identification of coproheme decarboxylase as a heme synthesis enzyme was a major step in understanding this newly appreciated pathway (13–15). However, the enzyme responsible for coproporphyrin production from coproporphyrinogen in the absence of oxygen has not been defined.

The anaerobic conversion of coproporphyrinogen to coproporphyrin is an outstanding question in coproporphyrin-dependent heme synthesis. In *S. aureus* and other organisms that synthesize heme via coproporphyrin, UroD produces coproporphyrinogen (Fig. 1A) from the common precursor uroporphyrinogen. Coproporphyrinogen oxidase (CgoX) has been identified as the next enzyme in the pathway (Fig. 1A). CgoX performs the oxidation of coproporphyrinogen III to coproporphyrin (Fig. 1B), and *in vitro* this reaction uses molecular oxygen as the electron acceptor (12, 16–18). This suggests the existence of an enzyme capable of producing coproporphyrin anaerobically, using a different electron acceptor. Existence of separate oxygen-dependent and -independent enzymes for this step in *S. aureus* would be in line with the oxygen-dependent and oxygen-independent synthesis of protoporphyrin IX in the protoporphyrin-dependent heme synthesis pathway (10). Based on literature and bioinformatic analyses, three different possibilities exist to resolve this gap in coproporphyrin-dependent heme synthesis: HemN (NWMN_1486) acts as an oxygen-independent coproporphyrinogen III dehydrogenase, the DUF1444 protein (NWMN_1636) acts as an oxygen-independent coproporphyrinogen III dehydrogenase, or CgoX functions both anaerobically and aerobically (Fig. 1B). NWMN_1486 is annotated as an oxygen-independent coproporphyrinogen III oxidase in bioinformatic databases, making it an obvious candidate for this enzymatic step because it contains a radical S-adenosyl-L-methionine motif, a HemN fold, and high amino acid conservation to the prototypical HemN proteins that perform the parallel step in the protoporphyrin-dependent heme pathway (19, 20). Recent evidence from other species suggests that this annotation is likely incorrect; based on sequence homology, NWMN_1486 likely belongs to a protein family which functions as heme chaperones (11, 21, 22). However, the function of NWMN_1486 has not been studied. On the other hand, NWMN_1636 is a protein of unknown function but has been proposed as the possible oxygen-independent coproporphyrinogen III oxidase based on genomic context (10). This suggestion was included in the findings of a large-scale analysis of the coproporphyrin-dependent pathways across *Firmicutes* and *Actinobacteria*; a protein with the DUF1444 domain was found to cooccur with terminal heme synthesis enzymes in *Firmicutes* (10). Therefore, we set out to test these possibilities and determine which enzyme produced coproporphyrin anaerobically.

CgoX is required for aerobic and anaerobic heme synthesis. To experimentally test the contribution of CgoX, NWMN_1486, and NWMN_1636 to heme synthesis, in-frame unmarked deletions were created in each corresponding gene, as well as in *uroD*. We selected *uroD* as a control; a mutant of this gene is unable to synthesize heme




FIG 1 CgoX is required for aerobic and anaerobic heme synthesis. (A) Overview of heme synthesis in *S. aureus*. (B) Details of the conversion of coproporphyrinogen to coproporphyrin performed aerobically by (Continued on next page)

but can still synthesize siroheme (Fig. 1A), the cofactor of nitrite reductase. *S. aureus* strains were grown in RPMI medium (without glucose), supplemented with Casamino Acids and glycerol as the primary carbon source, to quantitatively assess aerobic and anaerobic heme-dependent growth. As glycerol is nonfermentable, growth in this medium relies largely on oxygen-dependent (aerobic) or nitrate-dependent (anaerobic) respiration. *S. aureus* wild type (WT) grew robustly in this medium in the presence of oxygen, but the heme-deficient $\Delta cgoX$ and $\Delta uroD$ mutants were unable to grow (Fig. 1C; see also Fig. S1A in the supplemental material). The $\Delta NWMN_{1486}$ mutant grew robustly while the $\Delta NWMN_{1636}$ mutant had a modest growth defect. Under these conditions, addition of nitrate did not enhance growth as oxygen is the preferred terminal electron acceptor (Fig. 1D and Fig. S1B). Under anaerobic conditions, the $\Delta uroD$ mutant was unable to grow in the absence (Fig. 1E) or presence (Fig. 1F) of nitrate, consistent with heme being required for anaerobic growth under these conditions. Surprisingly the $\Delta cgoX$ mutant was also unable to grow when nitrate was provided, suggesting that CgoX is also required for anaerobic heme-dependent growth (Fig. 1E and F and Fig. S2). The $\Delta NWMN_{1486}$ and $\Delta NWMN_{1636}$ mutants were able to grow anaerobically when nitrate was provided, which is consistent with CgoX functioning aerobically and anaerobically, with no other gene being required for anaerobic conversion of coproporphyrinogen to coproporphyrin. As observed aerobically, the $\Delta NWMN_{1636}$ mutant had a modest growth defect anaerobically. Deletion of *cgoX* in the $\Delta NWMN_{1486}$ and $\Delta NWMN_{1636}$ mutant backgrounds reduced growth to the level of the $\Delta cgoX$ mutant alone. The growth of mutants lacking either *cgoX* or *uroD* was complemented by expressing the respective genes in *trans* (Fig. 1G and Fig. S3). Together, these data confirm that CgoX is required for anaerobic and aerobic growth under conditions which require heme and that $NWMN_{1486}$ and $NWMN_{1636}$ do not participate in anaerobic heme synthesis.

Anaerobic production of heme relies only on CgoX, which is functionally conserved among Firmicutes. CgoX appears to be the sole enzyme required for the conversion of coproporphyrinogen to coproporphyrin anaerobically, yet the $\Delta NWMN_{1636}$ mutant had a modest growth defect anaerobically. We therefore investigated whether $NWMN_{1486}$ or $NWMN_{1636}$ makes any contribution to anaerobic heme synthesis using the $\Delta cgoX \Delta NWMN_{1636}$ and $\Delta cgoX \Delta NWMN_{1486}$ mutants. Addition of 100 nM exogenous heme complemented the anaerobic growth of the $\Delta cgoX$, $\Delta uroD$, and $\Delta cgoX \Delta NWMN_{1486}$ mutants and partly complemented the growth of the $\Delta cgoX \Delta NWMN_{1636}$ mutant (Fig. 2A and Fig. S4A). We next assessed growth of the double mutants using a low concentration of heme to test whether $NWMN_{1486}$ or $NWMN_{1636}$ contributes to heme synthesis in the absence of *cgoX*. We chose 5 nM heme in case the 100 nM exogenous heme was in such excess that it masked any small effects. The growth of the $\Delta cgoX$ mutant was partly complemented by 5 nM exogenous heme, and the $\Delta cgoX \Delta NWMN_{1486}$ mutant grew indistinguishably when 5 nM heme was provided (Fig. 2B and Fig. S4B). These data suggest that $NWMN_{1486}$ has no role in anaerobic heme synthesis. Based on sequence homology, $NWMN_{1486}$ may be a heme chaperone (21) and does not appear to contribute to heme synthesis, but the contribution of this protein to *S. aureus* physiology remains to be explored. The $\Delta cgoX \Delta NWMN_{1636}$ mutant was partly complemented by 5 nM heme but still had a reduced growth rate compared to that of $\Delta cgoX$ mutant supplemented with heme. These data are consistent with the $\Delta NWMN_{1636}$ mutant having a general growth defect unrelated

FIG 1 Legend (Continued)

CgoX and potential enzymes functioning anaerobically. (C to F) Growth of *S. aureus* WT and indicated mutants in medium containing glycerol as the primary carbon source under the following conditions: aerobically (C), aerobically and supplemented with nitrate (D), anaerobically (E), and anaerobically and supplemented with nitrate (F). (G) Aerobic growth of WT and indicated mutants encoding plasmids with constitutive promoter or constitutive promoter upstream of *uroD* or *cgoX* in medium containing glycerol as the primary carbon source. OD₆₀₀, optical density at 600 nm.




FIG 2 Anaerobic production of heme relies only on CgoX, which is functionally conserved among *Firmicutes*. *S. aureus* WT and indicated mutants were grown in medium with glycerol as the primary carbon source and nitrate added as indicated. (A) Anaerobic growth with 100 nM exogenous heme added. (B) Anaerobic growth with 5 nM exogenous heme added. (C and D) Aerobic and anaerobic growth of the $\Delta cgoX$ strain complemented with *cgoX* of various species of *Firmicutes*. (E and F) Aerobic and anaerobic growth of the $\Delta cgoX$ strain complemented with *cgoX* of the *Actinobacteria* species *S. coelicolor*.

Downloaded from <http://msphere.asm.org/> on August 5, 2020 by guest

to heme synthesis. The growth defect of the $\Delta NWMN_{1636}$ mutant was complemented by providing *NWMN_{1636}* in *trans*, while exogenous heme had only a modest effect on its growth (Fig. S5). It is possible that *NWMN_{1636}* has some basal coproporphyrinogen III oxidase activity or somehow otherwise contributes to heme synthesis, as heme does increase the growth yield of the $\Delta NWMN_{1636}$ mutant. However, the absolute inability of the $\Delta cgoX$ mutant to grow aerobically or anaerobically (Fig. 1) suggests that any contribution of *NWMN_{1636}* to heme synthesis is minimal. Thus, the function of the DUF1444 family member *NWMN_{1636}* is still unknown.

Having established that CgoX functions in both aerobic and anaerobic heme-dependent growth, we tested whether the anaerobic function of CgoX was widespread among organisms that rely on coproporphyrin-dependent heme synthesis. The *cgoX* gene of the related *Firmicutes* species *Staphylococcus lugdunensis*, *Staphylococcus carnosus*, *Listeria monocytogenes*, and *Bacillus anthracis* complemented the aerobic and anaerobic growth defect of the *S. aureus* $\Delta cgoX$ mutant when provided in *trans* (Fig. 2C and D), suggesting that many members of the *Firmicutes* phylum use CgoX for aerobic and anaerobic heme synthesis. Many species of the phylum *Actinobacteria* also synthesize heme via the coproporphyrin intermediate. Interestingly, the *cgoX* gene of *Streptomyces coelicolor* was able to partly complement the aerobic growth of the *S. aureus* $\Delta cgoX$ mutant, suggesting that the enzyme is expressed but was unable to complement growth anaerobically (Fig. 2E and F). These data suggest that the anaerobic function of CgoX is not conserved across all organisms that synthesize heme via coproporphyrin. The inability of *S. coelicolor* *cgoX* to complement anaerobically is consistent with differences noted in the terminal steps of *Actinobacteria* heme synthesis. For instance, CpfC from *Firmicutes* does not require a cofactor while CpfC of *Actinobacteria* possess an iron-sulfur cluster, and ChdC in *Firmicutes* has a small lip near the active site absent in *Actinobacteria* (10). In conclusion, we have experimentally shown that CgoX is responsible for both aerobic and anaerobic heme synthesis in *S. aureus*, as well as in many heme-synthesizing species of the *Firmicutes* phylum. More work is needed to identify the electron acceptor CgoX uses anaerobically and to understand more fully the terminal steps of *Actinobacteria* heme synthesis.

Methods. (i) General growth and reagents. For bacterial strains, plasmids, and primers, see the supplemental material (Tables S1 to S3, respectively, and references 25 and 26). *S. aureus* strains were grown routinely on tryptic soy agar (TSA) or broth (TSB) supplemented with 10 $\mu\text{g/ml}$ chloramphenicol when necessary. When used, hemin chloride (referred to as heme) was used at concentrations noted in the text or on figures. Heme was prepared fresh at 10 mM in 0.1 M NaOH; for experiments in which heme was used, an equal volume of 0.1 M NaOH was used for all conditions. *Escherichia coli* strains were grown on lysogeny broth (LB) or LB agar (LBA), supplemented with 50 $\mu\text{g/ml}$ carbenicillin when necessary. For growth in aerobic liquid medium, an Innova44 incubator with shaking at 180 rpm was used. For standard cultures of 2 to 3 ml, 15-ml round-bottomed polypropylene tubes with aeration lids were used, at a 45° angle in the aerobic incubator or upright and without shaking in the anaerobic incubator. Unless noted otherwise, all chemicals are from Sigma. All molecular biology reagents were from New England Biolabs and used according to the manufacturer's instructions, unless otherwise noted. Phusion 2X high-fidelity master mix was used for all PCRs for cloning.

(ii) Gene deletions. In-frame, unmarked deletions were created by allelic exchange as described in Bae and Schneewind (23), with some modifications. The pKOR1 plasmids containing ~1-kb homologous regions flanking upstream and downstream of the gene to be deleted were prepared using NEB HiFi assembly according to manufacturer's suggestions. The pKOR1 backbone was amplified by PCR using primers JC291/JC292, which produce a linear product not including the *attB* recombination sites. The ~1-kb flanking regions were amplified from *S. aureus* Newman genomic DNA. During allelic exchange, 2 μM heme was added to the medium after generation of merodiploids to chemically complement heme synthesis defects. Deletions were confirmed by

PCR using isolated genomic DNA, and phenotypes were complemented by providing the gene in *trans*. For *uroD*, flanking regions were amplified using primers JC419/JC420 (upstream flanking) and JC421/JC422 (downstream flanking). For *cgoX*, flanking regions were amplified using primers JC631/JC632 (upstream flanking) and JC633/JC508 (downstream flanking). Deletions of *uroD* and *cgoX* were confirmed using primers JC427/JC428. For *NWMN_1636*, flanking regions were amplified using JC621/JC622 (upstream flanking) and JC623/JC624 (downstream flanking). The deletion was confirmed with PCR using primers JC627/JC628. For *NWMN_1486*, flanking regions were amplified using JC617/JC618 (upstream flanking) and JC619/JC620 (downstream flanking). The deletion was confirmed with PCR using primers JC625/JC626. To create $\Delta cgoX$ $\Delta NWMN_1636$, pKOR1-*cgoX* was transduced as described previously (24) using bacteriophage $\phi 85$ into $\Delta NWMN_1636$, and allelic exchange was performed. To create $\Delta cgoX$ $\Delta NWMN_1486$, pKOR1-*NWMN_1486* was transduced into $\Delta cgoX$, and allelic exchange was performed, with 2 μ M heme supplemented for all steps.

(iii) Genetic complementation. To complement the $\Delta NWMN_1636$ mutant phenotypes, *NWMN_1636* was cloned from *S. aureus* Newman genomic DNA using primers JC644/JC645 with homology to pOS1 P_{lgt} digested with NdeI and BamHI (NEB) and ligated using NEB HiFi assembly mix. To complement the $\Delta cgoX$ mutant phenotypes, *cgoX* was cloned from *S. aureus* Newman genomic DNA using primers JC515/JC516 to incorporate a C-terminal FLAG tag with homology to pOS1 P_{lgt} digested with XhoI and BamHI (NEB) and ligated using NEB HiFi assembly mix. The *cgoX* genes of other species were cloned in the same manner, except that nested PCR was used to accommodate the length of the 3' primer incorporating the FLAG tag sequence, as noted below. Genomic DNA was used as the template for *Firmicutes* species, while pET15b-*cgoX* (*S. coelicolor*) (provided by the Dailey lab) was used as the template for *S. coelicolor cgoX*. For *S. carnosus*, primers JC735/JC736 were used to amplify *cgoX* from the genome and became the template for JC735/JC737 to amplify the final product; for *S. lugdunensis*, primers JC738/JC739 were used to amplify *cgoX* from the genome and became the template for JC738/JC740 to amplify the final product; for *L. monocytogenes*, primers JC744/JC745 were used to amplify *cgoX* from the genome and became the template for JC744/JC746 to amplify the final product; for *B. anthracis*, primers JC747/JC748 were used to amplify *cgoX* from the genome and became the template for JC747/JC749 to amplify the final product. *S. coelicolor cgoX* did not require a nested PCR and was amplified with primers JC791/JC792. This plasmid was sequenced to confirm its accuracy. The complementation plasmids were confirmed by restriction digest after isolation from DH5 α following transformation, transformed into *S. aureus* RN4220 by electroporation, and transduced as described previously (24) using bacteriophage $\phi 85$.

(iv) Growth curves. Growth curves were performed using overnight cultures of biological triplicates or quadruplicates depending on the experiment. After overnight growth, 1 μ l of overnight culture was added to 199 μ l of RPMI medium (no glucose; Gibco) supplemented with 1% Casamino Acids, 0.04% glycerol, and 40 mM nitrate (from sodium nitrate; Fisher) in a 96-well round-bottomed plate (Costar), and growth was monitored in a BioTek plate reader with shaking at 37°C. Data were graphed using GraphPad Prism. Three independent experiments were performed (with the exception of data showing complementation with 100 nM heme, which was from a single representative experiment), the mean of the biological triplicates or quadruplicates was calculated, and the three means were graphed with the standard errors of the means shown for each figure panel, except for panels in which no error bars are displayed for ease of viewing. The same data, with errors indicated, are presented as supplemental figures. For aerobic growth curves, strains were streaked to TSA and grown aerobically for 24 h at 37°C. Single colonies were used (except for heme-deficient strains, for which a few colonies were used) to inoculate overnight cultures in 3 ml of TSB in aeration tubes and grown for 15 h with shaking at 180 rpm in an Innova44 incubator at 37°C. For anaerobic experiments, a Coy anaerobic chamber was used, filled with a mix of 90% nitrogen, 5% carbon dioxide, and 5% hydrogen gases, and hydrogen levels were

monitored to ensure a minimum of 2% hydrogen concentration. Palladium catalysts (Coy) were used to remove any residual oxygen by reaction with hydrogen. A Coy static incubator was maintained at 37°C. Solutions and plasticware were allowed to equilibrate for >24 h inside the glove box before use. For anaerobic samples, strains were streaked to TSA and grown aerobically for 24 h at 37°C. TSA plates were moved into the anaerobic chamber, and single colonies were used (except for heme-deficient strains, for which a few colonies were used) to inoculate overnight cultures in 2 ml of anaerobic TSB in aeration tubes and grown for 16 h anaerobically without shaking at 37°.

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/mSphere.00235-19>.

FIG S1, TIF file, 0.6 MB.

FIG S2, TIF file, 0.7 MB.

FIG S3, TIF file, 0.5 MB.

FIG S4, TIF file, 1.1 MB.

FIG S5, TIF file, 0.3 MB.

TABLE S1, DOCX file, 0.01 MB.

TABLE S2, DOCX file, 0.01 MB.

TABLE S3, DOCX file, 0.01 MB.

ACKNOWLEDGMENTS

We thank members of the Skaar laboratory for critical feedback on the manuscript. We thank Harry Dailey and R. Sophia Weerth (University of Georgia) for helpful feedback on the manuscript and for providing the *S. coelicolor cgoX* gene.

This work was supported by National Institutes of Health grants R01AI069233 (to E.P.S.), R01AI101171 (to E.P.S.), and F31AI126662 (to J.E.C.).

REFERENCES

- Tong SY, Davis JS, Eichenberger E, Holland TL, Fowler VG, Jr. 2015. *Staphylococcus aureus* infections: epidemiology, pathophysiology, clinical manifestations, and management. *Clin Microbiol Rev* 28:603–661. <https://doi.org/10.1128/CMR.00134-14>.
- Holmgren E, Hederstedt L, Rutberg L. 1979. Role of heme in synthesis and membrane binding of succinic dehydrogenase in *Bacillus subtilis*. *J Bacteriol* 138:377–382.
- Gaupp R, Schlag S, Liebecke M, Lalk M, Gotz F. 2010. Advantage of upregulation of succinate dehydrogenase in *Staphylococcus aureus* biofilms. *J Bacteriol* 192:2385–2394. <https://doi.org/10.1128/JB.01472-09>.
- Hammer ND, Schurig-Briccio LA, Gerdes SY, Gennis RB, Skaar EP. 2016. CtaM is required for menaquinol oxidase *aa₃* function in *Staphylococcus aureus*. *mBio* 7:e00823-16. <https://doi.org/10.1128/mBio.00823-16>.
- Hammer ND, Reniere ML, Cassat JE, Zhang Y, Hirsch AO, Hood MI, Skaar EP. 2013. Two heme-dependent terminal oxidases power *Staphylococcus aureus* organ-specific colonization of the vertebrate host. *mBio* 4:e00241-13. <https://doi.org/10.1128/mBio.00241-13>.
- Burke KA, Lascelles J. 1975. Nitrate reductase system in *Staphylococcus aureus* wild type and mutants. *J Bacteriol* 123:308–316.
- Chang JP, Lascelles J. 1963. Nitrate reductase in cell-free extracts of a haemin-requiring strain of *Staphylococcus aureus*. *Biochem J* 89:503–510. <https://doi.org/10.1042/bj0890503>.
- Proctor RA, von Eiff C, Kahl BC, Becker K, McNamara P, Herrmann M, Peters G. 2006. Small colony variants: a pathogenic form of bacteria that facilitates persistent and recurrent infections. *Nat Rev Microbiol* 4:295–305. <https://doi.org/10.1038/nrmicro1384>.
- Choby JE, Grunenwald CM, Celis AI, Gerdes SY, DuBois JL, Skaar EP. 2018. *Staphylococcus aureus* HemX modulates glutamyl-tRNA reductase abundance to regulate heme biosynthesis. *mBio* 9:e02287-17. <https://doi.org/10.1128/mBio.02287-17>.
- Dailey HA, Dailey TA, Gerdes S, Jahn D, Jahn M, O'Brian MR, Warren MJ. 2017. Prokaryotic heme biosynthesis: multiple pathways to a common essential product. *Microbiol Mol Biol Rev* 81:e00048-16. <https://doi.org/10.1128/MMBR.00048-16>.
- Dailey HA, Gerdes S, Dailey TA, Burch JS, Phillips JD. 2015. Noncanonical coproporphyrin-dependent bacterial heme biosynthesis pathway that does not use protoporphyrin. *Proc Natl Acad Sci U S A* 112:2210–2215. <https://doi.org/10.1073/pnas.1416285112>.
- Lobo SA, Scott A, Videira MA, Winpenny D, Gardner M, Palmer MJ, Schroeder S, Lawrence AD, Parkinson T, Warren MJ, Saraiva LM. 2015. *Staphylococcus aureus* haem biosynthesis: characterisation of the enzymes involved in final steps of the pathway. *Mol Microbiol* 97:472–487. <https://doi.org/10.1111/mmi.13041>.
- Dailey HA, Gerdes S. 2015. HemQ: an iron-coproporphyrin oxidative decarboxylase for protoheme synthesis in Firmicutes and Actinobacteria. *Arch Biochem Biophys* 574:27–35. <https://doi.org/10.1016/j.abb.2015.02.017>.
- Mayfield JA, Hammer ND, Kurker RC, Chen TK, Ojha S, Skaar EP, DuBois JL. 2013. The chlorite dismutase (HemQ) from *Staphylococcus aureus* has a redox-sensitive heme and is associated with the small colony variant phenotype. *J Biol Chem* 288:23488–23504. <https://doi.org/10.1074/jbc.M112.442335>.
- Celis AI, Gauss GH, Streit BR, Shisler K, Moraski GC, Rodgers KR, Lukat-Rodgers GS, Peters JW, DuBois JL. 2017. Structure-based mechanism for oxidative decarboxylation reactions mediated by amino acids and heme propionates in coproheme decarboxylase (HemQ). *J Am Chem Soc* 139:1900–1911. <https://doi.org/10.1021/jacs.6b11324>.
- Corrigall AV, Siziba KB, Maneli MH, Shephard EG, Ziman M, Dailey TA, Dailey HA, Kirsch RE, Meissner PN. 1998. Purification of and kinetic studies on a cloned protoporphyrinogen oxidase from the aerobic bacterium *Bacillus subtilis*. *Arch Biochem Biophys* 358:251–256. <https://doi.org/10.1006/abbi.1998.0834>.
- Hansson M, Gustafsson MC, Kannangara CG, Hederstedt L. 1997. Isolated *Bacillus subtilis* HemY has coproporphyrinogen III to coproporphyrin III

- oxidase activity. *Biochim Biophys Acta* 1340:97–104. [https://doi.org/10.1016/s0167-4838\(97\)00030-7](https://doi.org/10.1016/s0167-4838(97)00030-7).
18. Dailey TA, Meissner P, Dailey HA. 1994. Expression of a cloned protoporphyrinogen oxidase. *J Biol Chem* 269:813–815.
 19. KEGG. 2019. *Staphylococcus aureus* subsp. *aureus* Newman: NWMN_1486. Retrieved from KEGG https://www.genome.jp/dbget-bin/www_bget?sae:NWMN_1486 (accession no. T00557). Accessed 27 March 2019.
 20. NCBI. 2019. Oxygen-independent coproporphyrinogen III oxidase [*Staphylococcus aureus* subsp. *aureus* str. Newman]. Retrieved from the NCBI Protein Database <https://www.ncbi.nlm.nih.gov/protein/150374498> (accession no. BAF67758.1). Accessed 27 March 2019.
 21. Abicht HK, Martinez J, Layer G, Jahn D, Solioz M. 2012. *Lactococcus lactis* HemW (HemN) is a haem-binding protein with a putative role in haem trafficking. *Biochem J* 442:335–343. <https://doi.org/10.1042/BJ20111618>.
 22. Haskamp V, Karrie S, Mingers T, Barthels S, Alberge F, Magalon A, Muller K, Bill E, Lubitz W, Kleeberg K, Schweyen P, Broring M, Jahn M, Jahn D. 2018. The radical SAM protein HemW is a heme chaperone. *J Biol Chem* 293:2558–2572. <https://doi.org/10.1074/jbc.RA117.000229>.
 23. Bae T, Schneewind O. 2006. Allelic replacement in *Staphylococcus aureus* with inducible counter-selection. *Plasmid* 55:58–63. <https://doi.org/10.1016/j.plasmid.2005.05.005>.
 24. Choby JE, Mike LA, Mashruwala AA, Dutter BF, Dunman PM, Sulikowski GA, Boyd JM, Skaar EP. 2016. A small-molecule inhibitor of iron-sulfur cluster assembly uncovers a link between virulence regulation and metabolism in *Staphylococcus aureus*. *Cell Chem Biol* 23:1351–1361. <https://doi.org/10.1016/j.chembiol.2016.09.012>.
 25. Duthie ES, Lorenz LL. 1952. Staphylococcal coagulase: mode of action and antigenicity. *J Gen Microbiol* 6:95–107. <https://doi.org/10.1099/00221287-6-1-2-95>.
 26. Bubeck Wardenburg J, Williams WA, Missiakas D. 2006. Host defenses against *Staphylococcus aureus* infection require recognition of bacterial lipoproteins. *Proc Natl Acad Sci U S A* 103:13831–13836. <https://doi.org/10.1073/pnas.0603072103>.