

RELATIONSHIP BETWEEN DRS2 AND KES1 CONTROLLING
PROTEIN TRAFFICKING IN *Saccharomyces cerevisiae*

By

Baby-Periyanyaki Muthusamy

Dissertation

Submitted to the faculty of the
Graduate School of Vanderbilt University
in partial fulfillment of the requirements
for the Degree of

DOCTOR OF PHILOSOPHY

In

Biological Sciences

December, 2009

Nashville, Tennessee

Approved:

Professor Todd R. Graham

Professor Kendal S. Broadie

Professor Katherine Friedman

Professor Larry Swift

To my husband, Paramasivam Natarajan,
my child, Akilan Paramasivam
and my parents

ACKNOWLEDGEMENTS

I would like to express my deep and sincere gratitude to my advisor Dr. Todd R. Graham for his persistent guidance, teaching, discussion, and enormous encouragement during my graduate study. Dr. Graham's enthusiastic attitude towards science, insightful direction, and expansive knowledge of the field have inspired me. I would also like to thank my thesis committee members, Dr. Kendal Broadie, Dr. Kathy Friedman, Dr. Larry Swift, and Dr. Chris Hardy for donating their time and knowledge for my research studies and future scientific career.

I would also like to express my thanks towards past and present lab members for their assistance and friendship during my time in the Graham Lab, specifically Sophie Chen, Ke Liu, Jing Xiao, Xiaoming Zhou, Adrian Pineda, Kavitha Surendhran, Ryan Baldrige, Tessy Sebastian, Jake Randolph and others. I also appreciate past and present undergraduates who worked with me in the lab, especially Rohini Khatri and Jason Wen. I would like to acknowledge Dr. Chris Janetopoulos for access to his microscopy facilities.

I would like to thank funding agencies including National Institute of Health, Department of Biological Sciences and Graduate School of Vanderbilt University for research supports and research presentations in international conferences.

Lastly and most importantly I would like to thank my family members for their support without expectations. I would like to thank most specially to my husband Paramasivam Natarajan for his love, support, and encouragement in all aspects of my life. I want to thank my child Akilan Paramasivam for his energetic and curious outlook that inspires me during my study. I would like to thank my parents and my brother for their support and guidance and also both sides of my family for their encouragement in my life.

TABLE OF CONTENTS

	Page
DEDICATION.....	ii
ACKNOWLEDGEMENTS.....	iii
LIST OF TABLES.....	vii
LIST OF FIGURES.....	viii
 Chapter	
I. INTRODUCTION TO ROLE OF P4-ATPASES AND OXYSTEROL BINDING PROTEIN HOMOLOGUES IN PROTEIN TRANSPORT.....	1
P-type ATPases.....	2
Type IV P-type ATPases.....	10
A. P4-ATPases and membrane asymmetry.....	12
B. Functions of P4- ATPases in mammals.....	13
C. Roles of P4-ATPase in Yeast.....	18
D. Roles of P4-ATPase in other organisms.....	22
E. Evidence for lipid flippase function of P4-ATPases.....	23
1. Lipid flippase function for mammalian P4-ATPase.....	23
2. Lipid flippase function in yeast.....	24
Role of Drs2-Cdc50 in Protein Transport and Vesicle Biogenesis.....	26
Relationship of P4-ATPases to Sterols.....	34
A. Relationship between sterols and P4-ATPases in yeast and <i>C. elegans</i>	35
Proteins that bind Oxysterol.....	36
A. OSBP / ORP and their homologues.....	37
B. Osh proteins in Yeast.....	39
C. Osh4/Kes1.....	41
D. Evidence that Kes1 transports sterol.....	45
 II. CONTROL OF PROTEIN AND STEROL TRAFFICKING BY ANTAGONISTIC ACTIVITIES OF A P4-ATPASE AND OXYSTEROL BINDING PROTEIN HOMOLOGUE	 47

Introduction.....	47
Materials and Methods.....	51
Media and strains.....	51
Screen for suppressors of <i>drs2Δ</i> and cloning of <i>KES1</i>	53
Microscopy and Immunological Methods.....	57
Phospholipid translocase assays.....	57
Results.....	58
<i>kes1</i> suppresses <i>drs2Δ</i> cold-sensitive growth.....	58
<i>kes1Δ</i> suppression of <i>drs2Δ</i> requires <i>DNF</i> P4-ATPase genes.....	65
Kes1p attenuates Drs2p flippase activity at the TGN.....	69
Drs2p antagonizes the influence of Kes1p on sterol trafficking and sub- cellular distribution	74
<i>kes1Δ</i> does not suppress <i>drs2Δ</i> defects in protein trafficking between the TGN, plasma membrane and early endosomes.....	82
<i>kes1Δ</i> suppresses <i>drs2Δ</i> defects in TGN to vacuole protein transport.....	90
Discussion.....	95
 III. SUMMARY.....	 102
Cloning of <i>SDK</i>	103
Kes1 acts upstream to Drs2 in protein trafficking.....	107
PI4P in <i>kes1Δ</i>	107
Removal of Kes1 may improve the activity of Dnfs at the TGN.....	110
Kes1 may influence the sterol content of the TGN.....	112
Kes1 may influence the flippase activity by regulating a Protein phosphatase or kinase at the TGN.....	117
Future Directions.....	118
 REFERENCES.....	 119

LIST OF TABLES

Table	Page
1-1. The P-Type ATPase superfamily.....	3
1-2. The P4-ATPase in mammals.....	15
2-1. Yeast strain used in Chapter-II.....	54
2-2. Plasmids used In Chapter-II.....	55

LIST OF FIGURES

Figure	Page
1-1. Structure and catalytic cycle of P-type ATPase.....	6
1-2. P4-ATPase requirements for vesicle-mediated protein transport in the secretory and endocytic pathways.....	29
1-3. Structure of Kes1/Osh4.....	44
2-1. <i>kes1</i> is a suppressor of <i>drs2Δ</i> cold-sensitive and high-pressure growth defects.....	62
2-2. Whole cell lysates were prepared from strains carrying the indicated <i>KES1/SDK1</i> alleles and immunoblotted to detect Kes1p and glucose-6-phosphate dehydrogenase (G6PDH).....	63
2-3. Specificity for <i>kes1Δ</i> suppression of P4-ATPase mutants.....	68
2-4. Kes1p represses the flippase activity of Drs2p in TGN membranes.....	72
2-5. Relationship of ergosterol to <i>kes1Δ</i> suppression of <i>drs2Δ</i>	77
2-6. Enhanced rate of plasma membrane to ER sterol transport in <i>drs2Δ</i> cells is Kes1p dependent	81
2-7. <i>kes1Δ</i> does not suppress <i>drs2Δ</i> loss of plasma membrane phospholipids asymmetry.....	83
2-8. <i>kes1Δ</i> does not suppress <i>drs2Δ</i> defects in protein trafficking between the TGN, plasma membrane and early endosomes.....	86
2-9. <i>kes1Δ</i> does not suppress the <i>drs2Δ</i> defect in AP-1/clathrin- dependent trafficking of chitin synthase III between the TGN and early endosomes.....	89
2-10. <i>kes1Δ</i> suppresses <i>drs2Δ</i> defects in TGN to vacuole protein trafficking.....	93
3-1. Physical and Genetic interaction of Drs2 with traffic machinery proteins.....	106
3-2. Increased PI4P production in <i>kes1Δ</i>	109

3-3.	P4-ATPase requirements for vesicle-mediated protein transport in the secretory and endocytic pathways.....	111
3-4.	Potential role of Drs2 and Kes1 in establishing membrane curvature.....	114