

FACIAL EMOTION AND ORIENTATION-MATCHING IN SCHIZOPHRENIA PATIENTS
AND NORMAL CONTROLS

MIKISHA LYN DOOP

Thesis under the direction of Professor Sohee Park

Schizophrenia (SZ) patients exhibit deficits in recognition and identification of facial emotional expressions but it is unclear whether these deficits result from abnormal affective processing or an impaired ability to process complex visual stimuli such as faces. SZ and matched healthy controls (CO) participated in two computerized visual matching tasks (facial emotional expression and orientation). Accuracy and reaction time were recorded. Clinical symptoms were also assessed in the patients. SZ were less accurate on both facial emotion and orientation matching tasks compared with CO but there was no diagnosis-by-task interaction. Clinical symptoms of SZ were associated with deficits on the matching tasks and social functioning was correlated with emotion matching errors across both groups. SZ have general deficits in processing of faces, which is in turn associated with worse symptoms and reduced social functioning.

Approved _____ Date _____

FACIAL EMOTION AND ORIENTATION-MATCHING IN SCHIZOPHRENIA PATIENTS
AND NORMAL CONTROLS

By

Mikisha Lyn Doop

Thesis

Submitted to the Faculty of the
Graduate School of Vanderbilt University
in partial fulfillment of the requirements

for the degree of

MASTER OF ARTS

in

Psychology

August 2008

Nashville, Tennessee

Approved:

Sohee Park, Ph.D.

David Zald, Ph.D

To my father, James Albert Doop, and my mother, Frankie Ann Burchfiel, for learning to love
me no matter what.

To Professor Sohee Park, because plain old acknowledgements just won't cut it.

ACKNOWLEDGMENTS

This work would not have been possible without the financial support of the Vanderbilt Graduate School Fellowship and grants from NARSAD and NIMH. I am indebted to Dr. Andrew Tomarken, Chairman of the Department of Psychology, Dr. David Zald, and most importantly Dr. Sohee Park, who have been supportive of my career goals and who worked actively to provide me with the time to pursue those goals.

I am grateful to all of those with whom I have had the pleasure to work during this and other related projects. Both of the members of my Thesis Committee have provided me extensive personal and professional guidance and taught me a great deal about both scientific research and life in general. I would especially like to thank Dr. Sohee Park, the chairperson of my committee. As my teacher and mentor, she has taught me more than I could ever give her credit for here. She has shown me, by her example, what a good scientist (and person) should be.

I would also like to acknowledge Professor Tim Crow of Oxford University. His approach to scientific research renews my faith in the scientific process and gives me hope of a day where scientists will collaborate more and work together to find treatments, causes, and cures.

TABLE OF CONTENTS

	Page
DEDICATION	ii
ACKNOWLEDGEMENTS	iii
LIST OF TABLES	v
LIST OF FIGURES	vi
Chapter	
I. INTRODUCTION	1
II. METHODS	3
Participants	3
Materials and Procedures	5
Data Analysis	8
III. RESULTS	9
Accuracy	9
Reaction Time	10
Relationships with Social Functioning and Symptomatology	11
IV. DISCUSSION	13
REFERENCES	15

LIST OF TABLES

Table		Page
1.	Table 1: Demographic Data	4

LIST OF FIGURES

Figure		Page
1.	Figure 1: Emotion Matching Design	6
2.	Figure 2: Facial Orientation Matching Design	7
3.	Figure 3: Mean percent correct (bars are standard error)	9
4.	Figure 4: Mean reaction time in milliseconds (bars are standard error)	10

CHAPTER I

INTRODUCTION

Past research indicates that both vocal and facial emotion recognition is impaired in schizophrenia (Walker et al., 1984; Feinberg et al., 1986; Borod et al., 1993; Archer et al., 1994; Mandal et al., 1998; Edwards et al., 2001; Baudouin et al., 2002; Hooker & Park, 2002), and that this deficit is most likely a trait-like feature of the illness (Addington, et al., 2008; Exner et al., 2004; Kline et al., 1992; Poole et al., 2000; Salem et al., 1996; Schneider et al., 1995).

Schizophrenics (SZ) perform worse than control subjects (CO) on tasks that require emotion identification or discrimination (Walker et al., 1980; Salem et al., 1996) but such deficits may stem from a generalized deficit in face processing. Indeed evidence suggests that there is no specific deficit in emotion perception in SZ (Kerr & Neale, 1993) but that they may have generalized deficits in processing of faces (Salem et al., 1996) and face processing deficits may be related to cognitive deficits in schizophrenia (Bryson et al., 1997; Schneider et al., 1995; Addington & Addington, 1998; Kholer et al., 2000).

The relationships among symptom severity, emotion processing, and face processing have been investigated but are not clearly understood. Some studies have found a relationship between positive symptoms and face processing (Martin et al., 2005; Kholer et al., 2000; Baudouin et al., 2002; Schneider et al., 1995); other studies have found a relationship between negative symptoms and face processing (Suslow et al., 2003b; Mueser et al., 1996; Kholer et al., 2000), and yet other studies have found no relationship between symptoms and face processing

(Salem et al., 1996; Borod et al., 1993; Muzekari & Bates, 1977). Nevertheless, accurate processing of facial stimuli seems to have implications for social functioning in schizophrenia (Hooker & Park, 2002; Suslow et al., 2003a).

The goal of the present study was to extend previous findings of facial processing deficits of schizophrenia in relation to social functions using two simple matching tasks with no language or working memory demands which may have confounded past studies.

CHAPTER II

METHODS

Participants

Twenty-two healthy CO (12 females) were recruited via advertisements. CO were screened for history of psychiatric illness, head injury, epilepsy, and drug use. Sixteen (5 females) individuals who met the DSM-IV criteria for schizophrenia or schizoaffective disorder (13 schizophrenia, 3 schizoaffective) were recruited from an outpatient-clinic. All schizophrenic and schizoaffective subjects were chronically ill (mean years of illness = 13.4, S.D. = 7.4) and were out patient subjects. Exclusion criteria for SZ were multiple diagnoses, head injury, epilepsy, or current drug abuse. All SZ were taking atypical antipsychotics. Demographic information is presented in Table 1. All participants gave written informed consent approved by the Vanderbilt University Institutional Review Board and were compensated.

There were no group differences in age ($F(1,36) = 1.41, p = n.s.$), education ($F(1,36) = 0.41, p = n.s.$), estimated full-scale IQ ($F(1,34) = 0.92, p = n.s.$) and handedness measured by the Global Handedness Questionnaire ($F(1,36) = 1.96, p = n.s.$). There was a significant difference in Zigler Score of Social Functioning ($F(1,36) = 26.59, p < 0.0001$) between groups. IQ data were missing for 2 CO subjects but since the education level was matched, it is unlikely that the two groups differed in general cognitive ability.

Table 1: Demographic Data

	Control Subjects	Schizophrenic Subjects
	(n = 22)	(n = 16)
Education (years)	13.2 (1.9) ^A	12.8 (2.0)
Age	34.5 (11.2)	38.8 (10.2)
Full Scale I.Q.^B	95.4 (14.5)	100.0 (14.0)
BPRS	N/A	25.5 (13.7)
SANS	N/A	28.9 (19.3)
SAPS	N/A	26.3 (22.8)
SPQ	16.5 (8.0)	N/A
Global Handedness Score^C	80.3 (35.7)	58.3 (56.7)
CPZ Equivalent^D	N/A	283.9 (98.3)
Illness Duration	N/A	13.4 (7.4)
Zigler Score^E	5.0 (1.9)	2.2 (1.1)

^A Mean (standard deviation)

^BWechsler Adult Intelligence Scale - Revised (Wechsler, D., 1981) and Wechsler Abbreviated Scale of Intelligence (WASI; Wechsler et al., 1999).

^C Global Handedness Questionnaire (Ransil, B., & Schachter, S., 1994)

^D Chlorpromazine dose equivalent (in milligrams per day; Woods, 2003)

^EZigler Score of Social Functioning (Zigler, E., & Levine, J., 1981)

Materials and Procedures

All tasks were performed on an iMac computer (screen size 28.75 x 21.25cm). Subjects were seated 40 cm from the screen. Face stimuli were selected from the Karolinska Directed Emotional Faces (KDEF, Lundqvist et al., 1998). Calvo and Lundqvist (2008) conducted a study investigating valences and accuracy of identification using the KDEF stimuli. This study showed that happy faces were identified more accurately than other faces. However, fearful faces were identified less accurately than other faces. Please see www.psychonomic.org/archive for norms for each face and expression regarding identification accuracy, errors, and reaction times. Emotions used were neutral, happy, sad, fearful, and angry. All subjects had normal or corrected-to-normal vision. There were 20 trials per task and the order of task presentation was counterbalanced across subjects. For both tasks, accuracy and reaction times (RT) were recorded.

Emotion Matching

Subjects were asked to look at a fixation dot at the center of the screen. When they were ready to begin a trial, they clicked on the fixation dot with the mouse. The screen displayed a target face directly above the fixation dot and 3 different faces below the fixation dot (see Figure 1). Subjects were instructed to select from the face that matched the emotional expression of the target face above the fixation dot.




Figure 1: Emotion Matching Design

Facial Orientation-matching

The procedure was identical to the one described above for the emotion matching task but in this case, subjects were instructed to select the face that matched the orientation of the target face above the fixation dot (see Figure 2).




Figure 2: Facial Orientation Matching Design

Clinical symptoms

Brief Psychiatric Rating Scale (BPRS; Overall & Gorkhman, 1962), Scale for the Assessment of Positive Symptoms (SAPS; Andreasen, 1984), and Scale for the Assessment of Negative Symptoms (SANS; Andreasen, 1984) were used to assess symptoms.

Schizotypal Personality in Controls

We used the Schizotypal Personality Questionnaire (SPQ) (Raine, 1991), a self-report questionnaire based on DSM-III criteria for schizotypal personality disorder.

Social functioning

The Zigler social competence scale (Zigler & Levine, 1981) was used to estimate social functioning in all participants, using demographic information.

Data Analysis

Repeated measure ANOVA was used to compare the two groups on % correct responses and response times. Fisher's PLSD was used for the post-hoc analysis. Correlations were computed using the Pearson product moment correlation. Cohen's *d* was used to calculate effect sizes.

CHAPTER III

RESULTS

Accuracy

There was a main effect of diagnosis ($F(1,36)=10.10$; $p=0.003$, $r=0.89$). CO (Mean=92.50%; SD=8.46) performed better overall than SZ (Mean=81.41%; SD=20.25). A main effect for task was also found ($F(1,36)=8.82$; $p=0.005$, $r=0.84$). Subjects performed better on the facial orientation task (Mean=92.50%; SD=18.45) than on the emotion-matching task (Mean=83.16%; SD=10.16). There was no significant interaction (see Figure 3).




Figure 3: Mean percent correct (bars are standard error)

Reaction Time

There was no main effect of diagnosis ($F(1,36)=0.22$; $p=n.s.$), but there was a main effect of task type ($F(1,36)=20.09$; $p<0.0001$, $r=0.99$). Subjects were faster on the face-orientation task (Mean=4723ms; SD=3516) than on the emotion-matching task (Mean =7153ms; SD=2326).

There was a trend towards a group-by-task interaction ($F(1,36)=4.03$, $p<0.06$, $r=0.49$). However, post-hoc analyses showed no significant differences in reaction time between groups on either the facial orientation-matching task or the emotion-matching task (see figure 4).




Figure 4: Mean reaction time in milliseconds (bars are standard error)

Relationships with Social Functioning and Symptomatology

Across both groups, Zigler score of social function was positively correlated performance on the emotion-matching task ($r = 0.38$; $p=0.02$). Interestingly, there was also a trend for a significant correlation between Zigler score of social functioning and performance on the face-orientation task ($r=0.31$, $p=0.06$) suggesting that better performance on face processing in general may be related to better social functioning.

In SZ group, BPRS ratings were negatively correlated with performance on the emotion-matching task ($r = -0.57$, $p = 0.04$), suggesting that SZ with more symptoms were less accurate on the emotion-matching task. There was a trend towards a negative correlation between positive symptoms (SAPS) and accuracy on the emotion-matching task ($r = -0.52$, $p = 0.07$). There were no significant correlations with negative symptoms and task performance. Illness duration was negatively correlated with performance on the face-orientation task ($r = -0.68$, $p = .07$) and positively correlated with RT on the same task ($r = 0.74$, $p = 0.04$). Daily medication dose (as measured by the CPZ equivalent) was positively correlated with performance on the face-orientation task ($r = 0.53$, $p = 0.05$) and negatively correlated with reaction time on the same task ($r = -0.60$, $p = 0.02$). These findings suggest that symptom management is related to the ability to process facial stimuli. In other words, those patients with better face processing ability tend to have fewer symptoms.

CHAPTER IV

DISCUSSION

This study aimed to investigate face processing in SZ while controlling for external cognitive demands. Face processing was impaired in SZ whether they were asked to match emotion or orientation. It is interesting to note that while SZ tend to be slower than CO in general, there was no RT difference between SZ and CO in this study; both groups were faster on matching orientation than emotion.

Our data suggests that positive symptoms are related to poorer performance on emotion matching and may be related to a facial processing deficit in general. SZ patients with more symptoms performed worse on the emotion-matching task, and our data suggest that this finding may be a function of positive symptoms rather than negative symptoms. In line with this finding, Pinkham, et al. (2008) found that paranoid schizophrenics showed decreased neural activation in the right amygdala, fusiform face area, and left ventrolateral prefrontal cortex as compared to controls on a face processing task. These findings are interesting in that core features of negative symptoms are deficits of emotion processing, and one would hypothesize a relationship between negative symptoms and face processing deficits.

This study also underscores the relationship between social functioning and the ability to process affective facial stimuli. There was a positive correlation between the Zigler score of social functioning and the ability to correctly process emotional faces across both groups. This is similar to what Hooker and Park (2002) found in their study. They showed that affect

recognition performance was related to social functioning as measured by the Social Dysfunction Index (Hooker & Park, 2002).

There are several limitations to our study. The sample size is small so we were not able to examine the symptom-face processing relationship comprehensively. Secondly, it is yet to be determined if deficits in face processing are due to a larger overarching perceptual and cognitive deficit. Faces are after all, a class of complex visual stimuli. However, a study on biological motion perception in SZ patients (Kim et al, 2005) indicate that SZ may be selectively impaired in processing biological stimuli (i.e., socially-relevant stimuli such as faces) compared with nonbiological objects. Lastly, the Zigler score of social functioning is more equipped to measure pre-morbid social functioning than current social functioning. This may have influenced our findings when using that measure.

Taken together our findings indicate that SZ have deficits in face processing in general, replicating Salem et al's results (1996) and that face processing deficits may have significant socioaffective consequences for the patients.

REFERENCES

- Addington J, Penn D, Woods SW, Addington D, Perkins DO. 2008 Facial affect recognition in individuals at clinical high risk for psychosis. *British Journal of Psychiatry*. 192:67-8.
- Addington, J., Addington, D., 1998. Facial affect recognition and information processing in schizophrenia and bipolar disorder. *Schizophrenia Research* 32, 171–181.
- Andreasen, N.C., 1983. Scale for the Assessment of Negative Symptoms (SANS). University of Iowa, Iowa City.
- Andreasen, N.C., 1984. Scale for the Assessment of Positive Symptoms (SAPS). University of Iowa, Iowa City.
- Archer, J., Hay, D.C., Young, A.W., 1994. Movement, face processing, and schizophrenia: evidence of a differential deficit in expression analysis. *British Journal of Clinical Psychology* 33, 517–528.
- Baudouin, J.-Y., Martin, F., Tiberghien, G., Verlut, I., Frank, N., 2002. Selective attention to facial emotion and identity in schizophrenia. *Neuropsychologia* 40, 503–511.
- Borod, J.C., Martin, C.C., Alpert, M., Brozgold, A., Welkowitz, J., 1993. Perception of facial emotion in schizophrenia and right brain damaged patients. *Journal of Nervous and Mental Disease* 181, 494–502.
- Bryson, G., Bell, M., Lysaker, P., 1997. Affect recognition in schizophrenia: a function of global impairment or a specific cognitive deficit. *Psychiatry Research* 71, 105 – 113.
- Edwards, J., Pattison, P.E., Jackson, H.J., Wales, R.J., 2001. Facial affect and affective prosody recognition in first-episode schizophrenia. *Schizophrenia Research* 48, 235–253.
- Exner, C., Boucsein, K., Degner, D., Irle, E., Weniger, G., 2004. Impaired emotional learning and reduced amygdala size in schizophrenia: a 3-month follow-up. *Schizophrenia Research* 71, 493-503.
- Feinberg, T.E., Rifkin, A., Schaeffer, C., Walker, E., 1986. Facial discrimination and emotional recognition in schizophrenia and emotional disorders. *Archives of General Psychiatry* 43, 276–279.
- Hooker, C., Park, S., 2002. Emotion processing and its relationship to social functioning in schizophrenia patients. *Psychiatry Research*. 112, 41-50.
- Kerr, S.L., Neale, J.M., 1993. Emotion perception in schizophrenia: Specific deficit or further evidence of generalized poor performance? *Journal of Abnormal Psychology* 102, 312-318.

- Kim, J., Doop, M.L., Blake, R., Park, S., 2005. Impaired visual recognition of biological motion in schizophrenia. *Schizophrenia Research* 77, 299-307.
- Kline, J.S., Smith, J.E., Ellis, H.C., 1992. Paranoid and nonparanoid schizophrenic processing of facially displayed affect. *Journal of Psychiatric Research* 26, 169-182.
- Kohler, C.G., Bilker, W., Hagendoorn, M., Gur, R.E., Gur, R.C., 2000. Emotion recognition deficit in schizophrenia: association with symptomatology. *Biological Psychiatry* 48, 127-136.
- Lundqvist, D., Flykt, A., Ohman, A., 1998. *The Karolinska Directed. Emotional Faces*. Karolinska Institute: Stockholm.
- Mandal, M.K., Pandey, R., Prasad, A.B., 1998. Facial expressions of emotions and schizophrenia. *Schizophrenia Bulletin* 24, 399-412.
- Martin, F., Baudoin, J-Y., Tiberghien, G., Franck, N., 2005. Processing emotional expression and facial identity in schizophrenia. *Psychiatry Research* 134, 43-53.
- Mueser, K.T., Doonan, R., Penn, D.L., Blanchard, J.J., Bellack, A.S., Nishith, P., DeLeon, J., 1996. Emotion recognition and social competence in chronic schizophrenia. *Journal of Abnormal Psychology* 105, 271 - 275.
- Muzekari, L.H., Bates, M.E., 1977. Judgement of emotion among chronic schizophrenics. *Journal of Clinical Psychology* 32, 662-666.
- Overall, J.E., Gorhman, D.R., 1962. The brief psychiatric rating scale. *Psychological Reports* 10, 799-812.
- Pinkham, A.E., Hopfinger, J.B., Pelphrey, K.A., Piven, J., Penn, D.L. (2008) Neural bases for impaired social cognition in schizophrenia and autism spectrum disorders *Schizophrenia Research*. 99:164-75.
- Poole, J.H., Tobias, F., Vinogradov, S., 2000. The functional relevance of affect recognition errors in schizophrenia. *Journal of the International Neuropsychological Society* 6, 649-658.
- Raine, A., 1991. The SPQ: a scale for the assessment of schizotypal personality based on DSM-III-R criteria. *Schizophrenia Bulletin* 17, 555-564.
- Ransil, B.J., & Schachter, S.C. (1994). Test-retest reliability of the Edinburgh Handedness Inventory and Global Handedness preference measurements, and their correlation. *Perceptual and Motor Skills* 79, 1355-72.
- Salem, J.E., Kring, A.M., Kerr, S.L., 1996. More evidence for generalized poor performance in facial emotion perception in schizophrenia. *Journal of Abnormal Psychology* 105, 480-483.

Schneider, F., Gur, R.C., Gur, R.E., Shtatsel, D.L., 1995. Emotional processing in schizophrenia: neurobehavioral probes in relation to psychopathology. *Schizophrenia Research* 17, 67–75.

Spreen, O. and Strauss, E., 1998. A Compendium of Neuropsychological Tests: Administration, Norms, and Commentary (2nd Edition) Oxford University Press. Oxford.

Suslow, T., Roestel, C., Droste, T., Arolt, V., 2003a. Automatic processing of verbal emotion stimuli in schizophrenia. *Psychiatry Research* 120, 131-144.

Suslow, T., Roestel, C., Arolt, V., 2003b. Affective priming in schizophrenia with and without affective negative symptoms. *European Archives of Psychiatry and Clinical Neuroscience* 253, 292-300.

Walker, E., McGuire, M., Bettes, B., 1984a. Recognition and identification of facial stimuli by schizophrenics and patients with affective disorders. *British Journal of Clinical Psychology* 23, 37–44.

Walker, E.F., Marvit, S.J., Emory, E., 1980. A cross-sectional study of emotion recognition in schizophrenic. *Journal of Abnormal Psychology* 89:428-36.

Wechsler, D., 1981. *The Wechsler Adult Intelligent Scale Revised (WAIS-R)*. Harcourt Assessment, Inc., San Antonio, TX.

Wechsler, D., 1999. *Wechsler Abbreviated Scale of Intelligence (WASI)*. Harcourt Assessment, Inc., San Antonio, TX.

Woods, S.W. 2003. Chlorpromazine equivalent doses for the newer atypical antipsychotics. *Journal of Clinical Psychiatry* 64:663-7.

Zigler, E., Levine, J., 1981. Premorbid competence in schizophrenia: what is being measured? *Journal of Consulting and Clinical Psychology* 49, 96 – 105.